



TITLE:

Stable rank for a pair of C^* -algebras (C^* -algebras and its applications to topological dynamical systems)

AUTHOR(S):

Osaka, Hiroyuki

CITATION:

Osaka, Hiroyuki. Stable rank for a pair of C^* -algebras (C^* -algebras and its applications to topological dynamical systems). 数理解析研究所講究録 2000, 1151: 120-125

ISSUE DATE:

2000-04

URL:

<http://hdl.handle.net/2433/64072>

RIGHT:

Stable rank for a pair of C^* -algebras

立命館大学・理工 大坂 博幸 (Hiroyuki Osaka)

1 Introduction and Main Result

The (topological) stable rank of Rieffel[19] is noncommutative generalization of the dimension of a compact Hausdorff space. In fact, when X is a compact Hausdorff space, the stable rank of $C(X)$ is $\left\lceil \frac{\dim X}{2} \right\rceil + 1$, where $\dim X$ is covering dimension of X . Recall that a unital C^* -algebra A has stable rank n if for any element a_1, a_2, \dots, a_n and $\varepsilon > 0$ there exist b_1, b_2, \dots, b_n in A such that

$$\begin{aligned} (1) & \|a_i - b_i\| < \varepsilon \\ (2) & \sum_{i=1}^n b_i^* b_i > 0. \end{aligned}$$

The condition (2) is equivalent to that there exist c_1, c_2, \dots, c_n in A such that $\sum_{i=1}^n c_i b_i = 1$. If A has no unital, we define stable rank of A as stable rank of the unitization of A . Note that stable rank one condition is equivalent to that the set of invertible elements is dense in a given C^* -algebra.

Many mathematicians tried to determine stable rank of interesting C^* -algebras, in particular, simple unital C^* -algebras ([5] [6] [8] [10] [11] [12] [13] [14] [15] [18] [20] [21] [22] etc). For examples, AF C^* -algebras and non-commutative tori have stable rank one ([18]), Toeplitz algebra has stable rank two, and Cuntz algebra has an infinity ([19]).

It has been a problem of considerable interest to determine stable rank of a crossed product algebra $A \rtimes_\alpha G$ of a unital C^* -algebra A with stable rank one by a finite group G . Blackadar presented this problem in the case that A is an AF C^* -algebra ([2]), and constructed a symmetry α on $A = C[0, 1] \otimes UHF$ whose crossed product algebra $A \rtimes_\alpha \mathbb{Z}_2$ has stable rank two. So, to consider the above problem, we need the assumption of the simplicity on a given C^* -algebra A .

In this direction Jeong and the author conclude ([10][11]) that a crossed product algebra $A \rtimes_\alpha G$ has the cancellation property if A is simple with stable rank one and the SP-property. Recall that a C^* -algebra A is said to have the SP-property if any non-zero hereditary subalgebra of A has non-zero projection. For example, an AF C^* -algebra has the SP-property. Therefore, we could conclude by [1] that a crossed product algebra $A \rtimes_\alpha G$ has stable rank one if we add real rank zero condition to this crossed product algebra, that is, the set of self-adjoint elements with finite spectra in $A \rtimes_\alpha G$ is dense in the set of self-adjoint elements. As Elliott presented a crossed product algebra $UHF \rtimes_\alpha \mathbb{Z}_2$ with real

rank one, however, we can not always hope that a given crossed product algebra has real rank zero.

In this talk we try to estimate stable rank of a given unital C^* -algebra B by stable rank of a C^* -subalgebra A with common unit. In case that B is a crossed product algebra of A by a finite group G , $sr(B) \leq sr(A) \times |G|$ ([11]). More generally, we have the following result:

Theorem 1 *Let $1 \in A \subset B$ be unital C^* -algebras. Suppose that B is a finitely generated left A -module, that is, there are some n elements v_1, v_2, \dots, v_n in B such that $\sum_{i=1}^n Av_i = B$. Then, $sr(B) \leq sr(A) \times n$.*

2 Stable rank

We prove main theorem with using the technique of matrix algebras. To this end the following lemma is needed.

Lemma 2 (Spatial case of Rieffel[19]) *Let $n \in \mathbb{N}$.*

$$sr(M_n(A)) \leq sr(A).$$

Proof. We will give a sketch of the proof. Suppose that $sr(A) = m$. Take m elements T_1, \dots, T_m from $M_n(A)$. Set $S = (T_1, T_2, \dots, T_m)^t$ in $M_{nm, m}(A)$. Let $(a_1, a_2, \dots, a_{nm})$ be the first row in S . Since $sr(A) = m$, we may assume that there exist c_2, \dots, c_{m+1} such that

$$c_2 a_2 + c_3 a_3 + \dots + c_{m+1} a_{m+1} = 1 - a_1.$$

Consider

$$\begin{pmatrix} 1 & c_2 & \dots & c_{m+1} & 0 & \dots & 0 \\ & 1 & & & & & \\ & & \ddots & & & & \\ & & & \ddots & & & \\ & & & & \ddots & & \\ & & & & & \ddots & \\ & & & & & & 1 \end{pmatrix} S.$$

Then, the new first row is $(1, b_2, \dots, b_{nm})^t$. Doing the iteration there is an invertible matrix $R \in M_{nm}(A)$ such that $RS = \text{diag}(1, S')$, $S' \in M_{nm-1, n-1}$. By induction there is $U \in M_{n-1, nm-1}$ such that $US' = I_{n-1}$. Note that $\|R^{-1} \text{diag}(1, S') - S\|$ is small.

Write

$$\begin{aligned} R^{-1} \text{diag}(1, S') &= (S_1, \dots, S_m)^t \\ \text{diag}(1, U)R &= (U_1, \dots, U_m), \end{aligned}$$

where $S_1, \dots, S_m, U_1, \dots, U_m$ are in $M_n(A)$. Then, we have $\|T_i - S_i\|$ is small, and $\sum_{i=1}^m U_i S_i = I_n$. \square

Definition 3 Define

$$Lg_n(A) = \{(a_1, a_2, \dots, a_n) \in A^n \mid \sum_{i=1}^n Aa_i = A\}.$$

Then, $sr(A) \leq n$ if and only if $Lg_n(A)$ is dense in A^n .

Proof of Theorem 1.

We give only the proof of the case of $sr(A) = 1$ and $G = \mathbf{Z}_2$. That is, we will show that $sr(A \times_\alpha \mathbf{Z}_2) \leq 2$ for a unital C^* -algebra A . In general case we can guess it from the proof of Lemma 2.

Take $a_0 + a_1u, b_0 + b_1u$ in $A \times_\alpha \mathbf{Z}_2$, where u is a unitary implementing α . Let $\varepsilon > 0$ be given. Consider

$$\begin{pmatrix} a_0 + a_1u \\ b_0 + b_1u \end{pmatrix} = \begin{pmatrix} a_0 & a_1 \\ b_0 & b_1 \end{pmatrix} \begin{pmatrix} 1 \\ u \end{pmatrix}.$$

Since $sr(M_2(A)) = 1$ by Lemma 2, there exists an invertible element

$$\begin{pmatrix} c_0 & c_1 \\ d_0 & d_1 \end{pmatrix} \in M_2(A) \text{ such that}$$

$$\left\| \begin{pmatrix} a_0 & a_1 \\ b_0 & b_1 \end{pmatrix} - \begin{pmatrix} c_0 & c_1 \\ d_0 & d_1 \end{pmatrix} \right\| < \frac{\varepsilon}{2}.$$

Consider

$$\begin{pmatrix} c_0 & c_1 \\ d_0 & d_1 \end{pmatrix} \begin{pmatrix} 1 \\ u \end{pmatrix} = \begin{pmatrix} c_0 + c_1u \\ d_0 + d_1u \end{pmatrix}.$$

Then, $(c_0 + c_1u, d_0 + d_1u) \in Lg_2(A \times_\alpha \mathbf{Z}_2)$, and $\|a_0 + a_1u - (c_0 + c_1u)\| < \varepsilon$, $\|b_0 + b_1u - (d_0 + d_1u)\| < \varepsilon$. Hence, $sr(A \times_\alpha \mathbf{Z}_2) \leq 2$. \square

Corollary 4 Let $1 \in A \subset B$ be a pair of unital C^* -algebras, and $E : B \rightarrow A$ be a faithful conditional expectation of index-finite type. That is, there exists a quasi-basis $\{v_i^*, v_i\}_{i=1}^n$ such that $x = \sum_{i=1}^n E(xv_i^*)v_i$, $\forall x \in B$. Then, $sr(B) \leq sr(A) \times n$.

Corollary 5 Let $1 \in A$ be a unital C^* -algebra and G be a finite group. Then,

$$sr(A \times_\alpha G) \leq sr(A) \times |G|.$$

3 Application

Using Corollary 5 we can present an affirmative data to a question of Blackdar[2]:

Question 6 *Let A be a AF C^* -algebra and G be a finite group. Then*

$$sr(A \times_{\alpha} G) \leq 1.$$

Theorem 7 (Jeong-Osaka[11]) *Let A be a simple unital C^* -algebra with $sr(A) = 1$ and SP-property. If G is a finite group and α is an action of G on A then the crossed product $A \times_{\alpha} G$ has cancellation.*

Here, a C^* -algebra has SP-property if each of its non-zero hereditary C^* -subalgebras contains a non-zero projection.

In particular,

Corollary 8 *Under the assumptions of the above theorem, if $A \times_{\alpha} G$ has real rank zero, that is, any self-adjoint element can be approximated by a self-adjoint element with finite spectra, then $sr(A \times_{\alpha} G) = 1$.*

Remark 9 *Generally, we can not hope that a given simple crossed product algebra $A \times_{\alpha} G$ has real rank zero, even if A is UHF, and $G = \mathbf{Z}_2$ [7].*

If one consider a crossed product by the integer group \mathbf{Z} then there is no conditional expectation of index-finite type from the crossed product $A \times_{\alpha} \mathbf{Z}$ onto A , but we have the following cancellation theorem:

Theorem 10 (Jeong-Osaka[11]) *Let A be a simple unital C^* -algebra with $sr(A) = 1$ and SP-property. If α is an outer action of the integer group \mathbf{Z} on A such that $\alpha_* = id$ on $K_0(A)$ then the crossed product $A \times_{\alpha} \mathbf{Z}$ has cancellation.*

Example 11 *Simple AF C^* -algebras and non-commutative tori A_{θ} are examples for C^* -algebras in Theorems 7 and 10.*

参考文献

- [1] B. Blackadar, *Comparison Theory for simple C^* -algebras*, Operator algebras and Applications, LMS Lecture Notes, no. 135, Cambridge University Press, 1988.
- [2] B. Blackadar, *Symmetries of the CAR algebra*, Annals of mathematics, 131(1990), 589 - 623.
- [3] L. G. Brown, *Stable isomorphism of hereditary subalgebras of C^* -algebras*, Pacific J. Math. 71(1977), 335 - 348.

- [4] L. G. Brown and G. K. Pedersen, *C*-algebras of real rank zero*, J. Funct. Anal. 99(1991), 131 - 149.
- [5] L. G. Brown and G. K. Pedersen, *On the geometry of the unit ball of a C*-algebra*, J. reine angew. Math. 469(1995), 113 - 147.
- [6] M. Dadarlat, G. Nagy, A. Nemethi, and C. Pasnicu, *Reduction of topological stable rank in inductive limits of C*-algebras*, Pacific J. Math. 153(1992), 267 - 276.
- [7] G. A. Elliott, *A classification of certain simple C*-algebras*, Quantum and Non-Commutative Analysis (ed. H. Araki et al), 373 - 385, 1993 Kluwer Academic Publishers. Printed in the Netherlands.
- [8] N. E. Hassan, *Rangs stables de certaines extensions*, J. London Math. Soc. 52(1995), 605 - 624.
- [9] M. Izumi, Lecture at Tokyo Metropolitan University, 1997.
- [10] J. A Jeong and H. Osaka, *Extremally rich C*-crossed products and cancellation property*, J. Australian Math. Soc.(Series A) 64(1998), 285 - 301.
- [11] J. A Jeong and H. Osaka, *Stable rank of crossed products by finite groups*, submitted.
- [12] N. S. Larsen and H. Osaka, *Extremal richness of multiplier algebras and corona algebras of simple C*-algebras*, J. Operator Theory 38(1997), 131 - 149.
- [13] V. Nistor, *Stable range for tensor products of extensions of \mathbf{K} by $C(X)$* , J. Operator Theory 16(1986), 387 - 396.
- [14] V. Nistor, *Stable range for a certain class of type I C*-algebras*, J. Operator Theory 17(1986), 365 - 373.
- [15] H. Osaka, *Real rank of crossed products by connected compact groups*, Bull. London Math. Soc. 27(1995), 257 - 264.
- [16] H. Osaka, *SP-property for a pair of C*-algebras*, submitted.
- [17] M. Pimsner and S. Popa, *Entropy and index for subfactors*, Ann. Sci. Ecole Norm. Sup. (4) 19(1986), 57 - 106.
- [18] N. Riedel, *On the topological stable rank of irrational rotation C*-algebras*, J. Operator Theory 13(1985), 143 - 150.
- [19] M. A. Rieffel, *Dimension and stable rank in the K-theory of C*-algebras*, Proc. London Math. Soc. 46(1983), 301 - 333.

- [20] A. J-L. Sheu, *Cancellation theorem for projective modules over the group C^* -algebras of certain nilpotent Lie groups*, Canad. J. Math. 39(1987), 365 - 427.
- [21] T. Sudo and H. Takai, *Stable rank of the C^* -algebras of nilpotent Lie groups*, Internat. J. Math. 6(1995), 439 - 446.
- [22] T. Sudo and T. Takai, *Stable rank of the C^* -algebras of solvable Lie groups of type I*, J. Operator Theory 38(1997), 67 - 86.
- [23] Y. Watatani, *Index for C^* -algebras*, Memories of the Amer. Math. Soc. 424(1990).